

QCD at zero baryon density

Slavo Kratochvila^{a*} and Philippe de Forcrand^{a,b}

^aInstitute for Theoretical Physics, ETH Zürich, CH-8093 Zürich, Switzerland

^bCERN, Theory Division, CH-1211 Geneva 23, Switzerland

While the grand canonical partition function $Z_{GC}(\mu)$ with chemical potential μ explicitly breaks the Z_3 symmetry with the Dirac determinant, the canonical partition function at fixed baryon number $Z_C(B)$ is manifestly Z_3 -symmetric. We compare $Z_{GC}(\mu = 0)$ and $Z_C(B = 0)$ formally and by numerical simulations, in particular with respect to properties of the deconfinement transition. Differences between the two ensembles, for physical observables characterising the phase transition, vanish with increasing lattice size. We show numerically that the free energy density is the same for both ensembles in the thermodynamic limit.

1. INTRODUCTION

The grand canonical partition function with respect to the quark number

$$Z_{GC}(\mu) = \int [DU] e^{-S_g[U]} \det M(U; \mu), \quad (1)$$

which is commonly used, explicitly breaks the Z_3 -centre symmetry. We compare it with the canonical partition function at fixed baryon number B

$$Z_C(B) = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\bar{\mu}_I e^{-i3B\bar{\mu}_I} Z_{GC}(\mu = i\bar{\mu}_I T), \quad (2)$$

which preserves the Z_3 -center symmetry. Here, we consider the case $\mu = B = 0$ only. While in one ensemble the expectation value of the Polyakov loop is non-zero at any temperature, in the other it is always zero, as visible in Fig. 1. Therefore, an important question arises: Do both formulations agree in the thermodynamic limit for physically relevant observables? This is a subtle issue, leading to a controversy about contributions of non-zero *triality*¹ states. Refs.[1] suggest that non-zero triality (“unphysical”, fractional B) states give a non-vanishing contribution even in the thermodynamic limit, and must be taken into




Figure 1. Distribution of the complex Polyakov loop in the grand canonical (top) and canonical (bottom) ensembles. left: $4^3 \times 4$, right: $6^3 \times 4$. In the thermodynamic limit, the distributions agree for both ensembles, up to two additional Z_3 -rotations in the canonical ensemble.

account. In contrast, [2] shows, for QCD with staggered fermions, that the only relevant sector of the grand canonical $\mu = 0$ description in the thermodynamic limit is the $B = 0$ sector (the free energy densities of both descriptions are identical for $V \rightarrow \infty$). In other words, the canonical ensemble $Z_C(B = 0)$, which contains only zero-triality states, describes the same physics in the thermodynamic limit as the grand canonical ensemble $Z_{GC}(\mu = 0)$, which has contributions from all triality-sectors. Therefore, non-zero triality states do not contribute in this limit. By an explicit comparison of $Z_{GC}(\mu = 0)$ with $Z_C(B = 0)$, we hope to settle this discussion.

*Talk presented by S. Kratochvila

¹The difference of the quark and anti-quark numbers modulo 3, $t \equiv (\#Q - \#\bar{Q}) \bmod 3$, is called the quark *triality*.




Figure 2. Phase diagram of $Z_{GC}(i\mu_I)$ in the (μ_I, T) -plane. The transitions (full lines) are first order for four flavours of staggered quarks with mass $ma = 0.05$.

2. THE CANONICAL ENSEMBLE

We briefly recall the construction of the canonical partition function $Z_C(B)$. First, one fixes the number of quarks $N = \int d^3\vec{x} \bar{\psi}(\vec{x}) \gamma_0 \psi(\vec{x})$ to Q by inserting a δ -function in the grand canonical partition function:

$$Z_C(Q) = \int [DU][d\bar{\psi}][d\psi] e^{-S_g[U] - S_F[U, \bar{\psi}, \psi]} \delta(N - Q).$$

The δ -function admits a Fourier representation:

$$Z_C(Q) = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\bar{\mu}_I e^{-iQ\bar{\mu}_I} \times \int [DU][d\bar{\psi}][d\psi] e^{-S_g[U] - S_F[U, \bar{\psi}, \psi] + i\bar{\mu}_I N}$$

with the new variable $\bar{\mu}_I$. One recognizes $\mu_I = \bar{\mu}_I T$ as an imaginary chemical potential, so that

$$Z_C(Q) = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\bar{\mu}_I e^{-iQ\frac{\mu_I}{T}} Z_{GC}(i\mu_I).$$

The grand canonical partition function $Z_{GC}(i\mu_I)$ as a function of an imaginary chemical potential has interesting properties [3], see Fig. 2. It is even and $2\pi T/3$ -periodic in μ_I . For our purposes, the most important consequence is, that $Z_C(Q \neq 0 \bmod 3) = 0$. Therefore, we define the baryon number $B \equiv 3Q$ and end up with Eq.(2).

3. RESULTS

We simulate QCD with $N_f = 4$ flavours of staggered quarks of mass $ma = 0.05$, using Hybrid Monte Carlo. We measure the Polyakov loop, the

plaquette and the chiral condensate on lattices with extents 4^3 , 6^3 and $8^3 \times 4$ at seven different β 's, from $\beta_1 = 4.9$ to $\beta_7 = 5.1$. We analyse the results using multihistogram reweighting [4].

In order to sample the $B = 0$ canonical ensemble, we supplement the ordinary Hybrid Monte Carlo at fixed μ_I with a Metropolis update $\mu_I \rightarrow \mu'_I$, with acceptance $\min\left(1, \frac{\det^{N_f}(\mathcal{D}(\mu'_I) + m)}{\det^{N_f}(\mathcal{D}(\mu_I) + m)}\right)$. The determinant ratio is evaluated with a stochastic estimator, namely $\langle e^{-|\mathcal{D}(\mu'_I) + m|^{-N_f/2} (\mathcal{D}(\mu_I) + m)^{N_f/2} \eta^2 + |\eta|^2} \rangle_\eta$ where η is a gaussian complex vector. In addition, one can perform a “ Z_3 move” at any time (with acceptance 1): $\mu_I \rightarrow \mu_I \pm \frac{2\pi T}{3}$, $U_4(\vec{x}, t=0) \rightarrow U_4(\vec{x}, t=0) e^{\mp i \frac{2\pi}{3}}$, $\forall \vec{x}$.




Figure 3. Susceptibility of $\bar{\psi}\psi$ on all analysed lattices and both ensembles. left: $Z_{GC}(\mu = 0)$, right: $Z_C(B = 0)$. Even for the smallest, 4^4 , lattice, differences are barely visible.

Fig. 3 shows the susceptibility of the chiral condensate $\bar{\psi}\psi$ in both ensembles. The peak shifts slightly in β for the $4^3 \times 4$ lattices. For larger lattice sizes, the difference disappears and both ensembles indicate the same critical β_c already for the $8^3 \times 4$ -lattices. We observe the same behaviour for the average plaquette. This is our first evidence to answer the question about the influence of non-zero triality states: In the thermodynamic limit, non-zero triality sectors are suppressed and do not affect the results. The effect of non-zero triality sectors is smaller than the statistical errors.

In Fig. 4 we show a finite size analysis of observables in the two ensembles. We plot the minimum of the Binder cumulant $C_B(\mathcal{O}) = 1 - \frac{1}{3} \frac{\langle \mathcal{O}^4 \rangle}{\langle \mathcal{O}^2 \rangle^2}$ as a function of the inverse volume $1/V$. For both the average plaquette and the chiral condensate,




Figure 4. Binder cumulant minimum versus inverse volume for both ensembles (slightly shifted). left: average plaquette, right: chiral condensate. The thermodynamic extrapolation does not tend to $\frac{2}{3}$, indicating a first order transition.

the thermodynamic extrapolation does not tend to $\frac{2}{3}$ (top-limit of the plot) - indicative of a first order phase transition (as known from the literature [5]). Furthermore, the slopes of the linear fit are identical in both ensembles within statistical errors. Therefore, we cannot even claim smaller finite size effects in one of the two ensembles.




Figure 5. Free energy density as a function of μ_I . left side: $\beta = 4.95 < \beta_c$, right: $\beta = 5.10 > \beta_c$. In the latter case, a parabola nicely describes the data for the larger lattices.

In Fig. 5 we show the free energy density versus μ_I for both $\beta < \beta_c$ and $\beta > \beta_c$. In both cases, we clearly observe a minimum at $\mu_I = 0$. Therefore, in the thermodynamic limit, only $\mu_I = 0 \bmod \frac{2\pi T}{3}$ will survive. This establishes the equivalence of $Z_C(B = 0)$ with $Z_{GC}(\mu = 0)$. For $\beta = 4.95 < \beta_c$, we observe that the free energy density at $\mu_I = \pm\pi T/3$ seems to have zero derivative, indicating a crossover (as expected from the phase diagram Fig. 2). Instead, for $\beta = 5.10 > \beta_c$ we expect a cusp to develop, due to the first order phase transition. Indeed, it appears likely as the volume increases. A simple parabola gives a good description of the data, with a curvature in line with expectations from high temperature [6].

4. CONCLUSION

The thermodynamic limits of $Z_{GC}(\mu = 0)$ and $Z_C(B = 0)$ seem to agree with respect to the deconfinement phase transition. The effect of non-zero triality sectors is smaller than the statistical errors, already for $8^3 \times 4$ lattices. Therefore, these states can be included or excluded without affecting the results in the thermodynamic limit. Moreover, finite size effects are equivalent even on the smallest, $4^3 \times 4$, lattice.

The canonical formulation requires a centre-symmetric formulation of QCD, which can be achieved very simply with negligible computer overhead, by adding a single d.o.f. μ_I updated by Metropolis. Hence, the presence of a fermionic determinant does not necessarily break the centre-symmetry.

Note, that another grand canonical partition function $Z_{GC}(\mu)$ can be built from the $Z_C(B)$'s, which is completely centre-symmetric. This partition function will give identical expectation values to the usual $Z_{GC}(\mu)$, except for centre-sensitive observables (Polyakov loop), whose different behaviour can be understood by invoking spontaneous center-symmetry breaking [7].

REFERENCES

1. M. Oleszczuk and J. Polonyi, *Annals Phys.* **227** (1993) 76; M. Oleszczuk, *Nucl. Phys. Proc. Suppl.* **39BC** (1995) 471; O. Borisenko, M. Faber and G. Zinovev, *Mod. Phys. Lett. A* **12** (1997) 949, *Nucl. Phys. B* **444** (1995) 563.
2. V. Azcoiti and A. Galante, *Phys. Lett. B* **444** (1998) 421.
3. A. Roberge and N. Weiss, *Nucl. Phys. B* **275** (1986) 734.
4. A. M. Ferrenberg and R. H. Swendsen, *Phys. Rev. Lett.* **61** (1988) 2635.
5. M. Fukugita and A. Ukawa, *Phys. Rev. Lett.* **57** (1986) 503.
6. F. Karsch, K. Redlich and A. Tawfik, *arXiv:hep-ph/0306208*.
7. Ph. de Forcrand and S. Kratochvila, in writing.